

Research Statement

ASVIN G^{*†}

I have a broad interest in arithmetic geometry and related subjects and especially enjoy developing and applying tools from varied fields to questions in arithmetic geometry (and vice-versa). As some highlights of my interdisciplinary bend, [Andrew O'Desky](#) and I (Project [1.1](#)) develop a combinatorial framework to compute motivic invariants of certain natural moduli spaces in algebraic geometry. In the other direction, my ongoing work (Project [1.2](#) with [Colin Crowley](#), [Connor Simpson](#) and [Botong Wang](#)) tackles an outstanding conjecture on matroids, taking inspiration from algebro-geometric methods.

Within arithmetic geometry, a major theme of my work concerns moduli spaces. I have also recently developed an interest in arithmetic dynamics and I combine these two interests to study the dynamics of Hecke correspondences on moduli spaces and more specifically, "unlikely and just likely intersections" (Project [2](#)).

In the near future, I hope to study several topics in related areas. Most concretely, my ongoing Project [3.2](#) has a possible extension to motivic integration that we would like to pursue. I am also interested in extending the results of Project [3.1](#) to other settings - most notably the p -adic setting over \mathbb{F}_q (as opposed to the ℓ -adic setting in my paper) and non-abelian towers.

More speculatively, I am interested in the p -adic information to be gleaned from matroids. There is a general principle in arithmetic that for "integral" objects, the p -adic places often play just as important a role as the complex places. For instance, there are p -adic analogues of the classical Riemann zeta function and other such zeta functions and there has been much work in the past few decades in extending classical cohomology theories to ℓ -adic or p -adic ones. Despite matroids being integral objects with a purely combinatorial definition, most if not all of the recent work on the subject has concerned archimedean questions (such as log concavity for independent sets). I would like to explore p -adic properties of the various standard invariants constructed from matroids.

Another topic that I am interested in exploring is the role of p -adic geometry in arithmetic dynamics. There is a venerable tradition of tackling questions in this field using p -adic methods starting at least from the proof of the Skolem-Mahler-Lech theorem and continuing to this day, for instance in the work of Junyi Xie on the Zariski dense orbit conjecture [[Xie22](#)]. I hope to make further contributions along these lines leveraging my familiarity with p -adic arithmetic.

I give a brief synopsis of my various projects here and expand on them in the rest of this statement.

- **Project [1.1](#):** [[OG22](#)] is joint work with [Andrew O'Desky](#). It is concerned with the development of novel combinatorial tools in order to compute motivic invariants of moduli spaces of polynomials with a fixed factorization type.
- **Project [1.2](#):** This is a project in progress (joint with [Colin Crowley](#), [Connor Simpson](#) and [Botong Wang](#)). We aim to prove an equivariant version of various log concavity statements related to matroids.

* Department of Mathematics; University of Wisconsin, Madison.

† email: gasvinseeker94@gmail.com, website: asving.com

- **Project 2:** This project concerns the question of "unlikely and just likely intersections" on Shimura varieties. In [G.22b], I prove the first general results towards a conjecture on unlikely intersections of arbitrary subvarieties on Shimura varieties formulated by Shou Wu-Zhang's AIM group in the context of a product of modular curves. In [GHS22], Ananth N. Shankar, Qiao He and I prove the conjecture for Hilbert modular varieties over a finite field (with mild restrictions).
- **Project 3.1:** [G.22a] proves a novel result in non-abelian Iwasawa theory in order to prove a geometric statement about the convergence of certain characteristic polynomials in a tower of curves. At its core is an elementary statement regarding a matrix generalization of Fermat's little theorem.
- **Project 3.2:** In joint work with Yifan Wei and John Yin, we vastly generalize and prove a conjecture [BCFG22, Conjecture 1.2] about p-adic densities of polynomials with a given factorization type. In doing so, we prove a version of Chebotarev's theorem for non-archimedean local fields like \mathbb{Q}_p .

1 COMBINATORICS

1.1 Configuration spaces, graded spaces, and polysymmetric functions

Recall the following classical result of Gauss.

Theorem 1 (Gauss). *The number of monic, irreducible degree d polynomials in 1 variable over the finite field \mathbb{F}_q is given by*

$$N_{d,1}(q) = \frac{1}{d} \sum_{m|d} \mu(m) q^{d/m}.$$

One is naturally led to ask the same question when the polynomials are allowed to have many variables. In this context, one needs to make a distinction between *arithmetic* and *geometric* irreducibility¹. For technical reasons, we find it more useful to work with homogeneous polynomials in $n + 1$ variables and we define $N_{d,n}^{\text{arith}}(q)$ (respectively $N_{d,n}^{\text{geom}}(q)$) to be the number of arithmetic (respectively geometric) irreducible degree d polynomials in $n + 1$ variables up to scaling by \mathbb{F}_q^\times . We prove:

Theorem 2. *We have the following generalizations of Gauss' result.*

$$N_{d,n}^{\text{arith}}(q) = \frac{1}{d} \sum_{m|d} m \mu\left(\frac{d}{m}\right) \sum_{\lambda \vdash m} (-1)^{\ell-1} \frac{(\ell-1)!}{n_1! \cdots n_m!} \prod_{k=1}^m \left(1 + q + \cdots + q^{\binom{n+k}{k}-1}\right)^{n_k},$$

$$N_{d,n}^{\text{geom}}(q) = \frac{1}{d} \sum_{m|d} m \mu\left(\frac{d}{m}\right) \sum_{\lambda \vdash m} (-1)^{\ell-1} \frac{(\ell-1)!}{n_1! \cdots n_m!} \prod_{k=1}^m \left(1 + q^{d/m} + \cdots + q^{d/m[\binom{n+k}{k}-1]}\right)^{n_k}.$$

In proving this result, it is useful to consider the moduli space $M_{d,n}$ of irreducible homogeneous polynomials of degree d in $n + 1$ variables (up to scaling). It is an open subset of the projective space $X_{d,n} = \mathbb{P}^{\binom{n+d}{d}-1}$ and crucially, polynomial multiplication induces maps $X_{d,n} \times X_{e,n} \rightarrow X_{d+e,n}$ that makes the collection $\{X_{d,n} : n \geq 1\}$ into a graded monoid space. Whenever we have such a collection of spaces satisfying "unique factorization", [OG22, Theorem 0.1] computes the class of $M_{d,n}$ in terms of the class of $X_{d,n}$ in the Grothendieck ring of varieties which can be considered as a "universal" euler characteristic.

¹ i.e., irreducibility over \mathbb{F}_q versus irreducibility over $\overline{\mathbb{F}}_q$.

1.1.1 Polysymmetric functions and types

In order to prove these general results, we define $P\Lambda$, the ring of *polysymmetric functions* as an extension of the classical theory of symmetric functions. Very briefly, one can consider polysymmetric functions as an extension of the theory of symmetric functions where we allow weighted variables [OG22, Part 1]. We extend several constructions from the classical theory including a generalization of partitions called *types* which record the possible factorizations of a degree d polynomial.

This notion of a type is central to our work. They faithfully model factorizations of elements in any unique factorization domain and reflect many interesting details about expressing polynomials in a maximally factored form. Of most interest to us is the notion of an *arrangement number* $a(\tau, \lambda)$ between two types τ, λ . In short, this records the number of ways of rearranging a factorization of type τ into a factorization of type λ and can be used to define a partial order on types. This partial order (and an associated symmetry) are interesting even when restricted to just partitions.

We use the notion of types to give analogues of the classical elementary and homogeneous symmetric polynomials in $P\Lambda$. These bases are, respectively, a combinatorial reflection of the motives of the space of irreducible polynomials and the space of all polynomials.

1.1.2 Graded plethysm and the Grothendieck ring of varieties

The second ingredient that is needed is the notion of *graded plethysm*, extending the classical notion of plethysm on lambda rings. Lambda rings were originally defined by Grothendieck in the context of the Grothendieck-Riemann-Roch theorem. The prototypical example of a lambda ring is the ring of representations of a group and the lambda operations axiomatize the operations of symmetric and exterior products.

The Grothendieck ring of motives $K_0(\text{var})$ also has the structure of a lambda ring but it is deficient in one important respect - it is not a *special* lambda ring. We use the ring of polysymmetric functions $P\Lambda$ as a replacement for $K_0(\text{var})$ in order to prove our theorems.

In this process, we identify two especially interesting quotients of $K_0(\text{var})$: one where the Adams operations do act in a "special" way and a further quotient where they act trivially. The first one loses almost no information in terms of the usual motivic invariants while the second operation is quite a drastic one. The second one captures "Euler-characteristic" like motivic invariants and is much simpler to work with. It is plausible that we could completely understand the structure of this quotient of the Grothendieck ring.

1.1.3 Deeper strata

Types give a natural stratification of $X_{d,n}$ parametrizing the space of polynomials with a given factorization type. We give a recursive formula for these strata [OG22, Section 7]. In the quotient where the Adams operations act trivially, they can be computed in terms of the graded plethysm operations and admit fairly explicit formulas [OG22, Section 8].

1.2 Equivariant log concavity

There has been much recent work in proving the log concavity of sequences, inspired by techniques from algebraic geometry ([AHK18],[HW17] among many others). For a survey of the field, see [Huh22] and [Kal22].

To take a concrete example, consider a finite collection of vectors $\mathcal{V} \subset L^n$ in a vector space over a field L . For each positive integer k , we define $\mathcal{I}_k \subset \mathcal{P}(\mathcal{V})$ to be the collection of subsets of \mathcal{V} of size k consisting of linearly independent vectors. So for instance, \mathcal{I}_1 has elements corresponding to the non-zero elements in \mathcal{V} . If \mathcal{V} is itself a collection of linearly independent vectors, then \mathcal{I}_k is simply the

collection of all subsets of \mathcal{V} of size k . More generally, we can consider any *matroid* and consider the collection of independent sets of size k for each k as above.

Mason [Mas72] conjectured that the sequence $(\#\mathcal{I}_1, \#\mathcal{I}_2, \dots)$ is a log-concave sequence² for any matroid. This conjecture (and some stronger versions) have been recently proven by [AHK18],[ALGV18],[BH18],[BH20] among others. We consider a strengthening of this conjecture in a different direction. Returning to our example above, let G be a subgroup of the permutations of \mathcal{V} that preserve the notion of linear independence (and for an arbitrary matroid, we define G to be the automorphisms of the ground set that preserve the notion of independence). Then G will act on the sets \mathcal{I}_k and we define $P_k := k[\mathcal{I}_k]$ to be the corresponding permutation representations. One is naturally led to the following conjecture (Mason's conjecture corresponds to the case where G is the trivial group).

Conjecture 3. *For any $1 \leq a \leq b \leq c \leq d$ with $a + d = b + c$, there is a G equivariant injection $P_a \otimes P_d \rightarrow P_b \otimes P_c$.*

One can make analogous conjectures for various other algebraic invariants associated to a matroid such as the Orlik-Solomon algebra. [GPY17] and [MMPR21] formulate such conjectures, prove it in some very special cases and provide numerical evidence for the conjecture in general.

In our work so far, we have a refinement of the above family of conjectures where we define an *explicit* G -equivariant map $P_a \otimes P_d \rightarrow P_b \otimes P_c$ and it remains to prove that this is an injective map. If successful, our approach would give an alternate proof of Mason's conjecture bypassing the earlier work on this question.

2 UNLIKELY INTERSECTIONS OF SUBVARIETIES ON A SHIMURA VARIETY

There has been much recent interest on questions of *just likely and unlikely intersections* on Shimura varieties. As the simplest case, consider two families of elliptic curves \mathcal{E}, \mathcal{F} relative to C , a curve over a finite field or the ring of integers of a number field (possibly localized at some finite set of primes). In these cases, it is known under mild restrictions on the two families that there are infinitely many geometric points $x \in C$ such that the fibers \mathcal{E}_x and \mathcal{F}_x are isogenous to each other ([CO06, Proposition 7.3],[Cha14]).

To better understand this result, let us reformulate it in terms of moduli spaces. A family of elliptic curves over C gives rise to a map

$$\begin{aligned} C &\rightarrow X(1) \\ x &\rightarrow j(\mathcal{E}_x) \end{aligned}$$

where $X(1)$ is the modular curve so that our set up above corresponds to a map $f : C \rightarrow X(1) \times X(1)$. The geometric points $x \in C$ such that \mathcal{E}_x is isogenous to \mathcal{F}_x correspond exactly to the intersection of $f(C)$ with $T_N(\Delta)$, a Hecke translate of the diagonal $\Delta \subset X(1)^2$. Since $f(C)$ and $T_N(\Delta)$ have complimentary dimension, we expect there to be finitely many points in this intersection for any fixed N and the question is about the behaviour as $N \rightarrow \infty$. Such results have been generalized to families of abelian surfaces ([MST22b],[ST20]) and K3 surfaces ([MST22a],[SST19]).

Briefly, these papers consider the intersection of a curve C with a "special divisor" D in a Shimura variety and show that the number of distinct points in the intersection $C \cap T_N(D)$ grows without bound for $\{T_N : N \geq 1\}$ a sequence of Hecke translations. To do this, they bound the local intersection number $(C \cap T_N(D))_x$ uniformly in N but on the other hand, also show that the global intersection number $C \cap T_N(D) \rightarrow \infty$ as $N \rightarrow \infty$. These two facts together prove the desired result. Crucially, they provide the local bound by appealing to the moduli interpretation of the associated Shimura variety and applying deformation theoretic techniques while the global intersection number computation is related to the Fourier coefficients of a modular form using the "specialness" of D .

² i.e., $\#\mathcal{I}_k^2 \geq \#\mathcal{I}_{k-1}\#\mathcal{I}_{k+1}$.

Despite the proofs relying heavily on such moduli theoretic considerations, these problems can be naturally phrased for arbitrary subvarieties of a Shimura variety. Indeed, this was done by an AIM group working under Shou-Wu Zhang (Section 3.4).

Conjecture 4. *Let \mathcal{S} denote a simple Shimura variety (over some base), and suppose that V, W are generically ordinary subvarieties having complementary dimension. Then the set of points in V isogenous to some point of W is Zariski-dense in C . Further, the subset of $V \times W$ consisting of pairs of isogenous points is dense in $V \times W$.*

In this generality, the conjecture is known in very few cases. Perhaps most significantly, it is known in complete generality in the geometric case, i.e., over \mathbb{C} by [TT21]. The aforementioned work [Cha14] deals with the case where $\mathcal{S} = Y(1)$ relative to \mathcal{O}_k , the ring of integers of a number field (using the same strategy of proof as described before). In all other cases, it seems very hard to directly extend the older methods of proof to tackle the above conjecture when V, W are allowed to be arbitrary subvarieties.

We provide the first evidence for the above conjecture for arbitrary subvarieties in an *arithmetic* setting.

2.1 Families of totally split abelian varieties

In [G.22b, Theorem 6], I prove the above conjecture (suitably modified) for $\mathcal{S} = Y(1)^n$. This result naturally leads to the question where $\dim V + \dim W \ll \dim \mathcal{S}$. In the number field setting, the above is impossible if one of V, W is special by the Zilber-Pink conjecture (see [HP16, Conjecture 2.2]).

Over a finite field, [ST18] formulates the following heuristic in this direction. Let $\text{Is}_n(m)$ be the set of isogeny classes corresponding to the abelian varieties parameterized by $Y(1)^n(\mathbb{F}_{q^m})$. [DH98, Theorem 1.1] shows that the size of the set $\text{Is}_n(m)$ is about $q^{nm/2}$ (as $q^n \rightarrow \infty$). It seems reasonable to treat the map $i : Y(1)^n(\mathbb{F}_{q^m}) \rightarrow \text{Is}_n(m)$ sending a point $x \in Y(1)^n(\mathbb{F}_{q^m})$ to the isogeny class of the corresponding abelian variety A_x as a *random* map and suppose that every isogeny class is of size $q^{nm/2}$ (ignoring problems related to supersingularity and the like). Let $V, W \subset Y(1)^n$ now be curves. For $n \geq 2$ and m large enough, one would expect $|i(V(\mathbb{F}_{q^m}))| \sim q^m$ and $|i(V(\mathbb{F}_{q^m})) \cap i(W(\mathbb{F}_{q^m}))| \sim q^{2m}/q^{nm/2}$. Summing over all m , one expects that the number of points on V geometrically isogenous to a point on W is approximately $(1 - q^2/q^{n/2})^{-1}$ which is a finite quantity (for $n \geq 5$). Surprisingly, I find two infinite families of counter examples to this heuristic in [G.22b, Theorems 13,14] - there exist curves C_n, D_n in $X(1)^n$ (for n arbitrarily large) such that there are infinitely many points on C_n isogenous to some point on D_n .

2.2 Abelian surfaces with real multiplication

In [GHS22, Theorem 1.1], Qiao He, Ananth N. Shankar and I prove the conjecture for \mathcal{S} a Hilbert modular surface over \mathbb{F}_p with the restriction that p splits in the ring of endomorphisms. In this case, \mathcal{S} does not have a global product structure but we use our assumption on p , endomorphism ring to construct a *local* splitting of the Frobenius in order to complete the argument.

As a by-product of our analysis, we also compute the change of Faltings height under isogeny for Abelian surfaces parametrized by such \mathcal{S} .

3 p -ADIC ARITHMETIC

3.1 On the variation of the Frobenius in a non abelian Iwasawa tower

The eigenvalues of the Frobenius on the étale cohomology of a smooth, projective variety over a finite field carry significant arithmetic information. By the Weil conjectures, these eigenvalues are algebraic integers and their absolute values under any complex embedding are understood.

In [G.22a], I draw inspiration from Iwasawa theory to study the asymptotic behaviour of these eigenvalues in an "Iwasawa tower" and in particular, show that there is a strong ℓ -adic convergence statement to be made in many natural examples. Consider the following example.

Example 1. Consider the smooth projective curves C_n corresponding to the equations

$$Y^2 = X^{2^n} + 1 \text{ over } \mathbb{F}_5.$$

They define a tower $\cdots \rightarrow C_2 \rightarrow C_1$ with maps $C_{n+1} \rightarrow C_n$ defined by $(X, Y) \rightarrow (X^2, Y)$. The characteristic polynomial of σ_5 on $H_{\text{ét}}^1(\overline{C}_n, \mathbb{Z}_\ell)$ is

$$f_n(x) := \det \left(1 - \sigma_2 x | H_{\text{ét}}^1(\overline{C}_n, \mathbb{Z}_\ell) \right) = (1 - 2x + 5x^2) \prod_{i=1}^{n-2} (1 + x^{2^i} 5^{2^{i-1}})^2.$$

Note that $f_{n-1}(x)$ divides $f_n(x)$ and the inverse roots of $g_n(x) = f_n(x)/f_{n-1}(x)$ are of the form $\sqrt{5}\zeta$ for ζ a root of unity of order 2^{n-1} for $n \geq 3$.

In fact, I prove the same statement for towers of Fermat curves (from which the above follows) and Artin-Schreier curves. This prompts the question of what happens in a more general context. For instance, one could take a map $f : C \rightarrow \mathbb{P}^1$ or $f : C \rightarrow A$ for A an abelian variety of dimension d and pull back by the following diagrams:

$$\begin{array}{ccc} C_n & \xrightarrow{f_n} & \mathbb{P}^1 \\ \downarrow \pi_n & & \downarrow t \rightarrow t^{\ell^n} \\ C & \xrightarrow{f} & \mathbb{P}^1 \end{array} \quad \text{or} \quad \begin{array}{ccc} C_n & \xrightarrow{f_n} & A \\ \downarrow \pi_n & & \downarrow [\ell^n] \\ C & \xrightarrow{f} & A \end{array}$$

For simplicity of notation, we will only consider the first case here. Note that the $C_n \rightarrow C$ are geometrically (branched) Galois extensions with an abelian Galois group $G_n \cong (\mathbb{Z}/\ell^n \mathbb{Z})^b$ for $b = 1$ in Case A and moreover, the G_n themselves have an action of σ_q .

Let $M_n = H_{\text{ét}}^1(\overline{C}_n, \mathbb{Z}_\ell) / H_{\text{ét}}^1(\overline{C}, \mathbb{Z}_\ell)$ and define the characteristic polynomials

$$f_n(x) = \det (1 - \sigma_q x | M_n), g_n = \det (1 - \sigma_q x | M_n / M_{n-1}).$$

It does not seem to be true that g_n determines g_{n+1} as in Example 1. Nonetheless, the following weaker convergence statement is true. For simplicity again, we assume that q is a prime power such that $q - 1$ is divisible by ℓ but not ℓ^2 .

Theorem. [G.22a, Theorem 12] *There is a factorization*

$$f_m(x) = \prod_{n \leq m} h_n(x^{\ell^{n-1}})$$

for some polynomials $h_n(x) \in \mathbb{Z}[x]$ independent of m . Moreover, the following congruence

$$h_{n+1}(x) \equiv h_n(x) \pmod{\ell^n}$$

holds so that the limit $h_\infty(x) = \lim_n h_n(x)$ exists inside $\mathbb{Z}_\ell[x]$.

The convergence of the $h_n(x)$ is quite unexpected and striking and I prove it by proving a more general theorem about the convergence of certain sequences of matrices related to non-abelian Iwasawa theory. Note that this more abstract statement can be applied to many more geometric contexts than just our two examples of towers of curves above although I do not pursue this in our paper. It applies

to any tower of varieties with an action of an abelian group such that the Frobenius action on the cohomology has a "large" orbit. For instance, one could take hypersurfaces of the form

$$f(x_0^{\ell^n}, \dots, x_n^{\ell^n}) = 0 \subset \mathbb{P}_{\mathbb{F}_q}^n.$$

To keep notation simple, the following is a special (yet non-trivial) case of the general ℓ -adic convergence theorem.

Theorem. [G.22a, Theorems 7,8] *Let $F(t)$ be a $r \times r$ matrix with entries in $\mathbb{Z}_\ell[t]$. Suppose that q is a prime such that $q - 1$ is divisible by ℓ but not ℓ^2 . For each $n \geq 1$, define the matrix*

$$A_n = \prod_{i=1}^{\ell^n-1} F(\zeta_{\ell^n}^i)$$

with characteristic polynomial $p_n(x)$. Then, the limit $p_\infty(x) = \lim_n p_n(x)$ exists and more precisely, the following congruence is true:

$$p_{n+1}(x) \equiv p_n(x) \pmod{\ell^n}.$$

3.2 The density of polynomials with a given factorization type

Let us identify the points of \mathbb{P}^n over some field k with univariate polynomials (up-to scaling). Given a "factorization type" τ , we can form the subset of polynomials X_τ in $\mathbb{P}^n(k)$ with the given factorization type τ . For instance, τ might correspond to polynomials that factor completely over k . A natural question is to ask about the relative proportions of X_τ as we vary over τ . This question is highly sensitive to the base field - over an algebraically closed field all polynomials split completely while over a finite field \mathbb{F}_q , the limiting proportions (for large q) are given by the Chebotarev theorem.

There has been some recent work on this question where k is a non-archimedean local field such as \mathbb{Q}_p . In this setting, $\mathbb{P}^n(k)$ is naturally a topological space with a canonical probability measure (normalized so that the total measure is 1). In [BCFG22, Conjecture 1.2], they compute the "expected number of rational roots of a random polynomial over k " and observe that this number is a rational function in p and remarkably, is invariant under the transformation $p \rightarrow p^{-1}$. They conjecture that the density $\rho(\tau; p)$ of polynomials with a given factorization type τ is a rational function in p which also satisfies the remarkable symmetry $\rho(\tau; 1/p) = \rho(\tau; p)$.

In on-going work with Yifan Wei and John Yin, we generalize and prove the conjecture in the context of a finite, flat map $f : X \rightarrow Y$ that is generically Galois with group G between smooth, projective varieties (with some mild restrictions) over a non-archimedean local ring A with fraction field K . Given a $\text{Spec } K$ point of Y , one can associate to it the conjugacy class of a subgroup inside G coming from the functoriality of the étale fundamental group.

Conversely, if one fixed such a conjugacy class of a subgroup $H \subset G$, we can compute the density of K -points in Y realizing this conjugacy class. We prove that these densities satisfy the above invariance under a transformation generalizing the $p \rightarrow p^{-1}$ symmetry above.

REFERENCES

- [AHK18] Karim Adiprasito, June Huh, and Eric Katz. Hodge theory for combinatorial geometries. *Annals of Mathematics*, 188(2):381–452, 2018.
- [ALGV18] Nima Anari, Kuikui Liu, Shayan Oveis Gharan, and Cynthia Vinzant. Log-concave polynomials iii: Mason's ultra-log-concavity conjecture for independent sets of matroids. *arXiv preprint arXiv:1811.01600*, 2018.

- [BCFG22] Manjul Bhargava, John Cremona, Tom Fisher, and Stevan Gajović. The density of polynomials of degree n over \mathbb{Z}_p having exactly r roots in \mathbb{Q}_p . *Proceedings of the London Mathematical Society*, 2022.
- [BH18] Petter Brändén and June Huh. Hodge-riemann relations for Potts model partition functions. *arXiv preprint arXiv:1811.01696*, 2018.
- [BH20] Petter Brändén and June Huh. Lorentzian polynomials. *Annals of Mathematics*, 192(3):821–891, 2020.
- [Cha14] François Charles. Frobenius distribution for pairs of elliptic curves and exceptional isogenies. *arXiv preprint arXiv:1411.2914*, 2014.
- [CO06] Ching-Li Chai and Frans Oort. Hypersymmetric abelian varieties. *Pure Appl. Math. Q. Volume 2, Number 1, Special Issue: In honor of John H. Coates.*, pages 1–27, 2006.
- [DH98] Stephen A DiPippo and Everett W Howe. Real polynomials with all roots on the unit circle and abelian varieties over finite fields. *Journal of Number Theory*, 73(2):426–450, 1998.
- [G.22a] Asvin G. On the variation of the frobenius in a non abelian iwasaawa tower. *arXiv preprint arXiv:2203.16774*, 2022.
- [G.22b] Asvin G. Unlikely and just likely intersections for high dimensional families of elliptic curves. *arXiv preprint arXiv:2203.16420*, 2022.
- [GHS22] Asvin G., Qiao He, and Ananth N. Shankar. Just-likely intersections on hilbert modular surfaces. *arxiv:2209.02806*, 2022.
- [GPY17] Katie Gedeon, Nicholas Proudfoot, and Benjamin Young. The equivariant Kazhdan–Lusztig polynomial of a matroid. *Journal of Combinatorial Theory, Series A*, 150:267–294, 2017.
- [HP16] Philipp Habegger and Jonathan Pila. O-minimality and certain atypical intersections. In *Annales scientifiques de l’ENS*, volume 49. Société Mathématique de France, 2016.
- [Huh22] June Huh. Combinatorics and Hodge theory. 2022.
- [HW17] June Huh and Botong Wang. Enumeration of points, lines, planes, etc. *Acta Mathematica*, 218(2):297–317, 2017.
- [Kal22] Gil Kalai. The work of june huh. In *Proceedings of the International Congress of Mathematicians*, volume 28, 2022.
- [Mas72] John H Mason. Matroid: unimodal conjectures and Motzkin’s theorem. *Combinatorics*, 1972.
- [MMPR21] Jacob P Matherne, Dane Miyata, Nicholas Proudfoot, and Eric Ramos. Equivariant log concavity and representation stability. *arXiv preprint arXiv:2104.00715*, 2021.
- [MST22a] Davesh Maulik, Ananth N Shankar, and Yunqing Tang. Picard ranks of K_3 surfaces over function fields and the Hecke orbit conjecture. *Inventiones mathematicae*, pages 1–69, 2022.
- [MST22b] Davesh Maulik, Ananth N Shankar, and Yunqing Tang. Reductions of abelian surfaces over global function fields. *Compositio Mathematica*, 158(4):893–950, 2022.
- [OG22] Andrew O’Desky and Asvin G. Configuration spaces, graded spaces, and polysymmetric functions. *arXiv preprint arXiv:2207.04529*, 2022.

- [SST19] Ananth N Shankar, Arul Shankar, Yunqing Tang, and Salim Tayou. Exceptional jumps of Picard ranks of reductions of $K3$ surfaces over number fields. *arXiv preprint arXiv:1909.07473*, 2019.
- [ST18] Ananth N Shankar and Jacob Tsimerman. Unlikely intersections in finite characteristic. In *Forum of Mathematics, Sigma*, volume 6. Cambridge University Press, 2018.
- [ST20] Ananth N Shankar and Yunqing Tang. Exceptional splitting of reductions of abelian surfaces. *Duke Mathematical Journal*, 169(3):397–434, 2020.
- [TT21] Salim Tayou and Nicolas Tholozan. Equidistribution of Hodge loci ii. *arXiv preprint arXiv:2103.15717*, 2021.
- [Xie22] Junyi Xie. The existence of Zariski dense orbits for endomorphisms of projective surfaces. *Journal of the American Mathematical Society*, 2022.